
FAUNISTIC RECORDS FROM THE CZECH REPUBLIC – 268

Hymenoptera: Apocrita:**Chalcidoidea: Mymaridae**

Mymar taprobanicum Ward, 1875. Bohemia mer.: 4 km S of Veselí nad Lužnicí, Ruda Natural Reservation (7054), edge of bog, 49°09'3.611"N 14°41'30.593"E, 5.ix.2008, 1 ♀, yellow pan trap, P. Janšta lgt., det. et coll.; Moravia mer.: Hodonín env., Hurdis brickfield (7168), 48°52'25"N 17°09'07"E, 174 m a.s.l., 29.-30.vii.2008, 1 ♀, yellow pan trap, J. Farkač, J. Straka & V. Vrabec lgt., P. Janšta det. et coll. Apparently cosmopolitan species, known from Australia, Carribean, Palaearctic Region (Askew et al. 2001), India (Hayat 1992), Ivory Coast, Kenya, Madagascar, South Africa, USA, Costa Rica, Colombia, and New Zealand (Triapitsyn & Bereyovskiy 2001). New species for the Czech Republic.

Chalcidoidea: Torymidae

Idarnotorymus pulcher Masi, 1916. Moravia mer.: Rokytná env., Rokytnenské slepence National Nature Reserve, meadow above Tábor hill (6963), 49°03'37.85"N 16°19'45.121"E, vii.2005, 1 ♀, sweeping, P. Janšta lgt., det. et coll.; Oslavany, spoil heap of former Kukla mine (6864), 49°07'41"N 16°20'24"E, 271 m a.s.l., 28.-29.vii.2008, 1 ♀, yellow pan trap, J. Farkač, J. Straka & V. Vrabec lgt., P. Janšta det. et coll. Distributed in Bulgaria, Hungary (Grissell 1995), Croatia (Bouček 1977), Italy (Bouček 1965), Slovakia (Bouček 1956), Russia, Ukraine (Nikolskaya & Zerova 1978), and Uzbekistan (Bouček 1965). New species for the Czech Republic.

Apoidea: Crabronidae

Miscophus niger Dahlbom, 1844. Moravia mer.: Zbýšov env., spoil heap of former Jindřich mine (6864), 49°09'42"N 16°21'20"E, 371 m a.s.l., 28.-29.vii.2008, 1 ♂, yellow pan traps, J. Farkač, J. Straka & V. Vrabec lgt., J. Straka det. et coll. Distributed throughout Europe (Dollfuss 1991), occurrence in northern and eastern Europe scattered (Lomholdt 1975, 1976). In the Czech Republic known only from Bohemia (Straka & Farkač 2002). New species for Moravia.

Tachysphex tarsinus (Lepelletier, 1845). Bohemia bor.: Oleško env., Travčický les (5551), 6.v.2000, 1 ♀, P. Bogusch lgt., J. Straka det. et coll.; Moravia mer.: Hodonín env., Hurdis brickfield (7168), 48°52'25"N 17°09'07"E, 174 m a.s.l., 29.-30.vii.2008, 1 ♂, yellow pan traps, J. Farkač, J. Straka & V. Vrabec lgt., J. Straka det. et coll. Distributed in South and Central Europe, North Africa, Turkey, Kazakhstan, and China (Straka 2005); in Central Europe known from Austria, Germany, Hungary, Baltic coast of Poland (Bitsch et al. 2001), and Slovakia (Vepřek & Straka 2007). New species for the Czech Republic, found in both Bohemia and Moravia.